

RESEARCH

Open Access



Characteristics and recovery methods of studies falsely excluded during literature screening—a systematic review

Lisa Affengruber^{1,2*} , Andreea Dobrescu¹, Emma Persad¹, Irma Klerings¹, Gernot Wagner¹, Isolde Sommer¹ and Gerald Gartlehner^{1,3}

Abstract

Background: Due to the growing need to provide evidence syntheses under time constraints, researchers have begun focusing on the exploration of rapid review methods, which often employ single-reviewer literature screening. However, single-reviewer screening misses, on average, 13% of relevant studies, compared to 3% with dual-reviewer screening. Little guidance exists regarding methods to recover studies falsely excluded during literature screening. Likewise, it is unclear whether specific study characteristics can predict an increased risk of false exclusion. This systematic review aimed to identify supplementary search methods that can be used to recover studies falsely excluded during literature screening. Moreover, it strove to identify study-level predictors that indicate an elevated risk of false exclusions of studies during literature screening.

Methods: We performed literature searches for eligible studies in MEDLINE, Science Citation Index Expanded, Social Sciences Citation Index, Current Contents Connect, Embase, Epistemonikos.org, and Information Science & Technology Abstracts from 1999 to June 23, 2020. We searched for gray literature, checked reference lists, and conducted hand searches in two relevant journals and similar article searches current to January 28, 2021. Two investigators independently screened the literature; one investigator performed the data extraction, and a second investigator checked for correctness and completeness. Two reviewers assessed the risk of bias of eligible studies. We synthesized the results narratively.

Results: Three method studies, two with a case-study design and one with a case-series design, met the inclusion criteria. One study reported that all falsely excluded publications (8%) could be recovered through reference list checking compared to other supplementary search methods. No included methods study analyzed the impact of recovered studies on conclusions or meta-analyses. Two studies reported that up to 8% of studies were falsely excluded due to uninformative titles and abstracts, and one study showed that 11% of non-English studies were falsely excluded.

Conclusions: Due to the limited evidence based on two case studies and one case series, we can draw no firm conclusion about the most reliable and most valid method to recover studies falsely excluded during literature screening or about the characteristics that might predict a higher risk of false exclusion.

Systematic review registration: <https://osf.io/v2pjr/>

*Correspondence: lisa.affengruber@donau-uni.ac.at

¹ Department for Evidence-Based Medicine and Evaluation, Cochrane Austria, Danube University Krems, Dr. Karl Dorrek Strasse 30, 3500 Krems, Austria
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords: Single screening, Falsely excluded studies, Systematic review, Rapid review

Background

Screening of titles, abstracts, and full-text publications to identify potentially eligible studies is an essential methodological element of any evidence synthesis. To reduce the risk of erroneously excluding relevant studies during literature screening, most international evidence synthesis organizations recommend dual-reviewer screening, that is, having two persons independently screen titles, abstracts, and full texts [1–3].

Dual-reviewer screening, however, is time-consuming. Due to the growing need to provide evidence syntheses under time constraints, researchers have begun focusing on the exploration of rapid review methods, which streamline the methodological steps of the systematic review process to provide answers more quickly. Rapid reviews often use single-reviewer screening, whereby each record is screened by only a single investigator. Single-reviewer screening reduces screening time by approximately 60% when compared to dual-reviewer screening [4]. The downside of single-reviewer screening, however, is that it is prone to falsely excluding relevant studies. In a crowd-based randomized controlled trial, Gartlehner et al. reported that single-reviewer abstract screening missed on average 13% (sensitivity: 86.6%; 95% confidence interval [CI], 80.6 to 91.2%) of relevant studies [5]. By comparison, dual-reviewer screening missed an average of 3% (sensitivity: 97.5%; 95% CI, 95.1 to 98.8%) of relevant studies [5]. In a recent systematic review by Waffenschmidt et al. on single- versus dual-reviewer screening, the median proportion of relevant but missed studies with single-reviewer screening was 5% (range 0 to 58%) [6].

Consequently, methods to mitigate the risk of erroneous exclusions of relevant studies and to recover falsely excluded studies during the evidence synthesis process are critical to ensure the validity of rapid review results. Table 1 lists several methods that could be used in an

attempt to recover falsely excluded studies [7, 8]. These methods are based on supplementary literature searches that can identify studies with a high likelihood of being relevant to the topic of interest. For example, commonly employed methods include reference list checking of the included studies or of other systematic reviews or using the “similar articles” function in electronic databases. The principle behind these supplementary searches is that investigators can reconsider the inclusion or exclusion of articles with a high likelihood of being relevant. In some cases, they might identify studies that investigators falsely excluded during single-reviewer literature screening. However, to date, only half of the published rapid reviews conducted reference list checking of the eligible studies [9].

Additionally, it would be of interest to researchers to know whether certain types of studies or publications have a higher risk of being falsely excluded than others [10]. For example, the publication year could play a role, as abstract reporting standards have changed over time. Older studies might be falsely excluded more frequently during abstract screening since information now considered relevant for an abstract might not have been deemed as important at the time of publication. As another example, one explanation for the false exclusion of studies in the systematic review by Waffenschmidt et al. was that the research question was too vague and largely depended on the interpretation of the reviewer [6].

The aim of this study was to systematically assess which methods have been used to recover studies falsely excluded during literature screening. Additionally, we aimed to identify the potential predictors and characteristics of falsely excluded studies.

Methods

The aim of this systematic review was to address the following key questions (KQs):

Table 1 Definitions of commonly used methods that could recover falsely excluded studies

Reference list checking (backward citation tracking)	Checking the reference lists of the included studies and any relevant systematic reviews identified [7, 8]
Similarity searches (i.e., related articles)	Using a key article to identify additional relevant articles by using a “similar articles” option available in some databases and search engines (e.g., PubMed, Google Scholar) [7, 8]
Forward citation tracking of included studies	Using citation indexes for forward citation searching based on a key article [8]
Academic search engines	Keyword searches using an academic search engine (e.g., Google Scholar) [8]
Contacting experts/researchers/companies/other stakeholders	Contacting individuals and organizations for information about relevant studies [8]

KQ 1: How effective are different methods in recovering studies falsely excluded during literature screening?

KQ 2a: What are the characteristics of studies that have been falsely excluded during literature screening?

KQ 2b: Can predictors help identify studies that are at a high risk of being falsely excluded?

This systematic review was conducted according to Cochrane methods [11]. We followed the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) 2020 statement [12]. The PRISMA checklist can be seen in Additional file 1. We registered our study protocol “Falsely excluded studies in the literature screening process—a systematic review” at <https://osf.io> (<https://osf.io/5zdpb/>). Because we could not identify studies that formally assessed predictors of the false exclusion of records during literature screening, we amended the protocol on June 11, 2021. We expanded the inclusion criteria to also include the study characteristics of falsely excluded studies, even if these characteristics were not formally assessed in a predictive model.

Eligibility criteria

The a priori–defined eligibility criteria are listed in Table 2 and described in more detail below.

We searched for studies assessing the use of supplementary search methods (e.g., forward citation tracking, reference list checking, and web searching) to recover studies falsely excluded during literature screening. These supplementary search methods are defined in Table 1.

Additionally, we searched for studies focusing on the predictors and characteristics of falsely excluded studies, such as those based on study design or publication type. Based on internal discussion and consensus among co-authors, we generated a list of potential predictors of false exclusion. This list was not exhaustive, and any other predictor not named on the list would have been eligible. For detailed eligibility criteria, see Table 2.

Information sources

An experienced information specialist performed searches for eligible studies in MEDLINE (Ovid), Science Citation Index Expanded, Social Sciences Citation Index, Current Contents Connect (all via Web of Science), Embase (Elsevier), Epistemonikos.org, and Information Science & Technology Abstracts

Table 2 Inclusion and exclusion criteria

	Inclusion	Exclusion
KQ 1: Methods to recover falsely excluded studies	<ul style="list-style-type: none"> Reference list checking of included studies Similarity searches (i.e., related articles) Forward citation tracking of included studies using citation indexes Google Scholar search Contacting experts/researchers/companies/other stakeholders 	Other methods
KQ 2: Potential predictors of false exclusion	<ul style="list-style-type: none"> Study design Main objective Sample size Country of conduct Year of publication Structure and content of abstract (e.g., only title available, no abstract, uninformative abstract) Language of publication Risk of bias Database indexing of studies (e.g., PubMed listing) Publication type Journal in which the study is published Impact factor of journal in which the study is published Others 	
Outcomes	<ul style="list-style-type: none"> Proportion of falsely excluded studies that could be recovered (KQ 1) Recall, precision, numbers needed to read (NNR) of supplementary searches (KQ 1) Impact of recovered studies on meta-analysis results and conclusions (KQ 1) Falsely excluded studies by characteristics (KQ 2) Falsely excluded studies by predictors (KQ 2) 	Other outcomes
Study design	<ul style="list-style-type: none"> Systematic reviews (KQ 1, 2) Randomized/nonrandomized trials (KQ 1, 2) Prospective and retrospective, controlled and uncontrolled observational studies (KQ 1, 2) 	Nonempirical publications (e.g., editorials, letters)
Date of search	Published 1999 or later	1998 and earlier
Publication language	No restrictions	No restrictions

Abbreviations: KQ key question, NNR numbers needed to read

(Ebsco) from 1999 to June 23, 2020. We first developed a search strategy for Ovid MEDLINE and then amended it to fit other electronic databases. We considered publications in all languages. According to the peer review of the electronic search strategy (PRESS) statement [12], the electronic Ovid MEDLINE search strategy was peer-reviewed by another information specialist. See Additional file 2 for the database search strategies.

In addition, we searched for gray literature (i.e., unpublished studies) relevant to this systematic review. Potential sources of gray literature included the Open Science Framework (www.osf.io), websites of known organizations that produce rapid reviews (e.g., Canadian Agency for Drugs and Technologies in Health [CADTH]) based on the CADTH Gray Matters Checklist [13], and dissertation databases (e.g., Digital Access to Research Theses [DART]-Europe). Furthermore, we searched for Cochrane Colloquium abstracts of oral, poster, and workshop presentations and Health Technology Assessment international (HTAi) meeting abstracts.

We manually searched the reference lists of background articles on this topic for any relevant citations that our electronic searches might have missed. Additionally, we hand searched journals that regularly publish methods studies, such as *Systematic Reviews* and *Research Synthesis Methods*. If our search retrieved conference abstracts of studies that might have fulfilled our inclusion criteria, we manually searched for further information about these studies (e.g., publications, entries in trial registries, etc.). Additionally, an information specialist conducted similar articles searches for identified key articles in PubMed and Google Scholar and forward citation tracking using Scopus up to January 28, 2021. The search results for the similar article searches are ranked by “similarity” to the key article; the top 20 articles are those categorized as the most similar according to the search algorithm. We exported the top 20 articles and assessed them according to our eligibility criteria. See Additional file 3 for the similar articles searches and forward citation tracking.

Study records

Data management

Identified citations were stored in an EndNote® X8.2 bibliographic database (Thomson Reuters, New York, NY). All results of the abstract and full-text review, including the exclusion reasons during the full-text review, were recorded in the EndNote database. PDF files of all full-text articles were stored on a server accessible to all members of the review team.

Selection process

Deduplication of the search results was carried out with EndNote® X8.2 (Thomson Reuters, New York, NY). We developed and pilot-tested abstract and full-text review forms that reflected our inclusion and exclusion criteria. Two independent reviewers screened abstracts and full-text articles in Covidence (www.covidence.org) and evaluated their eligibility for inclusion. Any discrepancies were resolved through discussion or consultation with a third reviewer. A total of 50 abstracts were piloted by all reviewers to resolve discrepancies and to test the abstract review form. The full-text review form was piloted with five full-text articles.

Data collection process

We designed and pilot-tested a structured data abstraction form. The data were extracted by one reviewer and checked for completeness and accuracy by a second investigator. The data extraction process was piloted with five studies.

Data items

For studies that met our inclusion criteria, we extracted the following study characteristics and outcomes:

- Study characteristics: author, year of publication, aims, study design, sample size (e.g., number of studies analyzed), number of reviewers involved
- Characteristics of methods/information sources used to recover falsely excluded studies (for KQ 1)
- Characteristics of falsely excluded studies/publications: study design, content of the abstract, language of publication (for KQ 2)
- Outcomes: proportion of falsely excluded studies/publications that could be recovered, impact of recovered studies on meta-analysis results and/or conclusions, proportion of falsely excluded studies/publications by characteristic or predictor

Risk of bias assessment

For methods studies with a case-study design, we adapted the Joanna Briggs Institute Critical Appraisal Checklist for Case Reports, and for method studies with a case-series design, the Joanna Briggs Institute Critical Appraisal Checklist for Case Series [14].

Data synthesis

We summarized the results narratively and grouped them by outcomes of interest. We did not identify enough studies with a similar design to be able to conduct meta-analyses.

Results

The literature searches identified 3750 deduplicated unique records, of which 124 full texts were assessed for eligibility. Three studies published in four publications met our inclusion criteria [10, 15–17]. One study reported on KQ 1, all three included studies considered KQ 2a, and no study was identified for KQ 2b. Figure 1 depicts the record review flow. Additional file 4 lists the studies excluded at the full-text level and the reasons for exclusion.

Characteristics of the included studies

We included three studies (published in four articles), of which two were method studies with a case-study design [10, 16, 17], and one was a methods study with a case series design [15]. We rated the risk of bias of two studies as high [15, 17] and of one study as low [16]. Additional file 5 presents the detailed risk of bias ratings. The aim of the three studies was mainly to assess different methods to accelerate the literature screening process, but they also reported the study characteristics of falsely excluded studies and recovery methods. Rathbone et al. [15] and Feehan et al. [10, 17] mentioned

uninformative abstract content details as a study characteristic of the falsely excluded studies. The study by Busse et al. [16] reported falsely excluded studies by publication language, comparing English-speaking reviewers with native-speaking reviewers. The study by Feehan et al. additionally reported their methods to recover those studies falsely excluded during literature screening [10, 17].

For the study by Feehan et al. [17], we identified a companion publication. This conference abstract by Beck et al. [18] was mentioned in a Cochrane review [10] on “Checking reference lists to find additional studies for systematic reviews.” All studies added relevant information for evidence synthesis. Table 3 presents the characteristics of the included studies.

Three methods studies [10, 15–17] assessed studies falsely excluded during abstract screening; one study assessed only full-text screening [16]. The three studies [10, 15–17] assessed one to ten systematic reviews including 53 to 211 studies and involved five to 16 reviewers in their literature screening process. The results for KQs 1 and 2 are described in detail in the following sections and in Table 3.




Fig. 1 Flow diagram

Table 3 Characteristics and results of the included studies

Author, year	Key question	Study design	Risk of bias	Aim	N of analyzed SRs, studies, publications	Abstract and/or full-text screening	N of reviewers involved	Proportion of falsely excluded studies/publications by characteristic	Proportion of falsely excluded studies that could be recovered
Busse 2014 [16]	KQ 2	Methods study with a case-study design	Low	To explore the extent to which English-speaking reviewers can differentiate eligible from ineligible foreign-language articles in a systematic review of all treatments for fibromyalgia	1 SR, 53 studies	Full-text screening	16	Single screening: NR Dual screening: <i>Non-English language</i> : Overall: 6/53 (11.3%) German: 2/16 (12.5%) French: 1/4 (25.0%) Turkish: 1/9 (11.1%) Chinese: 1/7 (14.3%) Korean: 1/1 (100%) Italian: 0/5 (0%) Spanish: 0/5 (0%) Portuguese: 0/3 (0%) Dutch: 0/1 (0%) Russian: 0/1 (0%) Polish: 0/1 (0%)	NR
Feehan 2011 [10, 17]	KQ 1, KQ 2	Methods study with a case-study design	High	To track eligible articles identified by checking references lists and to determine which of those had also been found through database searches, but had been screened out by the review authors	1 SR, 119 studies, 134 publications	Abstract screening	8	Single screening: NR Dual screening: Abstract content details: 11/134 publications (8.2%) due to uninformative titles and abstracts of the bibliographic record	11/11 publications (100%) through reference list checking through contacting key authors, reference list checking, forward citation tracking

Table 3 (continued)

Author, year	Key question	Study design	Risk of bias	Aim	N of analyzed SRs, studies, publications	Abstract and/or full-text screening	N of reviewers involved	Proportion of falsely excluded studies/publications by characteristic	Proportion of falsely excluded studies that could be recovered
Rathbone 2017 [15]	KQ 2	Methods study with a case-series design	High	To evaluate the feasibility of PICO-based title-only screening for scoping searches and rapid reviews by measuring the reduction in screening effort and the maintenance of recall of relevant records	10 SRs, 211 studies	Abstract screening	5	Single screening: <i>Abstract content details:</i> 1/211 (0.005%) due to an uninformative title	Dual screening: NR

Abbreviations: KQ key question, N number, NA not applicable, NR not reported, PICO Participants, Intervention, and Comparator, but not the Outcome, SR systematic review

Methods to recover studies falsely excluded during literature screening (KQ 1)

One methods study with a case study design [10, 17] conducted a scoping review on an orthopedic topic and analyzed all references included in their review. We rated this study as having a high risk of bias due to it missing conclusions and clear descriptions of the characteristics of falsely excluded studies. The authors found that 11 of 134 eligible studies had been initially found by database searches but had been erroneously excluded during dual screening of the titles and abstracts. The investigators recovered all 11 falsely excluded references (100%) through reference list checking of the included studies [10, 17]. The investigators also performed other supplementary search methods, such as contacting key authors, reference list checking, forward citation tracking, and hand searches of online journal websites, but these methods did not recover any falsely excluded references. The study did not analyze the impact of recovered studies on conclusions or meta-analyses.

Characteristics or predictors of falsely excluded studies (KQ 2)

All three studies explored the characteristics of studies falsely excluded during literature screening. We could not identify studies that formally assessed whether specific characteristics could predict an increased risk of a study being falsely excluded.

The characteristics mentioned in the studies were uninformative abstract content details and a non-English publication language (Table 3). The following sections summarize these findings in more detail.

Abstract content details

Two methods studies, one with a case-study design and one with a case-series design, mentioned uninformative abstract content details as a characteristic for false exclusion [10, 15, 17]. We rated these two methods studies as having a high risk of bias mainly due to them analyzing a convenience case or case series and missing reports of the eligibility criteria and characteristics of the included cases. Feehan et al. [10, 17] documented false exclusions during the production of a scoping review including 134 publications. Eleven of the 134 publications (8%) were falsely excluded due to uninformative titles and abstracts [10, 17]. Rathbone et al. [15] assessed a convenience sample of 10 systematic review datasets including 211 eligible studies, derived from the literature searches of completed systematic reviews, to test Participants, Intervention, and Comparator, but not the Outcome (PICo)-based title-only screening. Only a single study was falsely excluded by PICo-based title-only single screening in one of the 10 systematic reviews. The authors reported that

“ventilation” was used in the title as an alternative term for oxygen therapy, and this was not listed in the MeSH (Medical Subject Headings) database nor found while searching other resources, and therefore, subject knowledge was needed to identify the study. They concluded that if authors use uncommon or ambiguous terminology in the abstract, this might lead to false exclusion [15].

Non-English publication language

One methods study with a case-study design, rated as low risk of bias, mentioned a publication language other than English as a characteristic for false exclusion during full-text screening. Busse et al. [16] evaluated a 10-question guide for English reviewers to assess the inclusion of non-English articles compared to native-language speakers. The authors reported false exclusion during the production of one systematic review of randomized controlled trials of fibromyalgia therapy including 53 publications published in 11 languages other than English. Six of the 53 full texts (11%) were falsely excluded by English-language reviewers due to being published in German (2/53), French, Turkish, Chinese, and Korean (1/53, respectively) languages [16].

Discussion

To the best of our knowledge, this is the first systematic review that addresses methods to recover falsely excluded studies and study characteristics that can potentially predict a higher risk of false exclusions during literature screening. Our review identified three studies [10, 15–17]. One study reported on methods to recover studies falsely excluded during literature screening (KQ 1), all three included studies considered the characteristics of falsely excluded studies (KQ 2a), and no study was identified on predictors that could help identify studies at a high risk of being falsely excluded (KQ 2b). For KQ 1, only one methods study with a case study design that evaluated supplementary search methods was identified. Only reference list checking recovered 100% of the studies missed by dual literature screening; other supplementary search methods, such as contacting key authors, forward citation tracking, and hand searches of online journal websites, did not recover any falsely excluded studies. The study did not analyze the impact of recovered studies on conclusions or meta-analyses. [10, 17]. Three studies [10, 15–17] reported on the characteristics of falsely excluded studies. Two studies [10, 15, 17] reported false exclusions of up to 8% of studies due to uninformative titles and abstracts. Another study [16] reported false exclusion of 11% of studies due to the non-English publication language.

For rapid reviewers, the results underline the importance of reference list checking as well as considering

uninformative titles/abstracts and non-English publications. It seems prudent that rapid review teams should check the reference lists of relevant publications to recover possible missed studies. Other supplementary search methods, such as contacting key authors, forward citation tracking, and hand searches of online journal websites, do not seem to be as effective. However, this evidence is derived from only a single study with a high risk of bias. Review team leaders should consider discussing uninformative titles and abstracts with screeners to avoid such false exclusions, perhaps by requesting that screeners include studies with uninformative titles or abstracts by default for a more in-depth evaluation at the full-text level. Journals and authors should strive to follow abstract reporting guidelines within the theme they are writing, considering the structure and content details of their abstract to ensure their studies are not falsely excluded. Journal editors and peer reviewers should also emphasize informative titles and abstracts. Additionally, the results show that authors should involve translation software or translators in the conduct of rapid and systematic reviews including non-English articles, either internally or through a research network. Generally, the growing use of automation software may prevent the false exclusion of studies [19]. This would require providing a clean, high-quality initial data set to train the algorithm, ensuring that any duplicates with conflicting decisions are removed and prespecifying records correctly as includes or excludes [19].

Overall, the quality and quantity of the included studies limit the evidence base of our systematic review. We identified three single-method studies on individual cases, assessing only 12 systematic reviews and 18 falsely excluded studies, that mostly address KQ 2. For KQ 1, we were only able to identify one method study with a case-study design. For KQ 2, we could only identify studies assessing the characteristics “Abstract content details” and “non-English publication language.” We were not able to identify studies addressing other characteristics or predictors of falsely excluded studies. We could not identify studies including predictive models for falsely excluded studies. It might be possible that reviewers falsely excluded studies due to other reasons (e.g., loss of concentration, reviewer’s experience) that were not formally addressed by this systematic review. Additionally, the review includes two high risks of bias studies of three included studies. We rated these two studies as having a high risk of bias mainly due to them analyzing a convenience case or case series and missing reports of the eligibility criteria and characteristics of the included cases. These flaws could have an influence on the reliability of our results. Therefore, the results should be cautiously taken into consideration.

Our systematic review has some methodological limitations. Although we applied a rigorous methodology according to Cochrane methods [11], we cannot rule out the possibility that we might have missed relevant studies. Another known threat to the validity of systematic reviews that we cannot exclude is publication bias. Although we searched for gray literature, relevant research on this topic might not be published due to non-significant results.

Based on our findings, there is an immense need for future research to evaluate supplementary search methods to recover studies falsely excluded during literature screening, particularly given the increased demand for rapid reviews employing literature screening shortcuts. No guidance document of any institution has addressed this issue. Studies assessing the overall impact of retrieving missed studies on conclusions and meta-analyses are also warranted. It is further unclear if falsely excluded studies are at risk of remaining permanently excluded. For example, a falsely excluded study may be retrieved again by reference list checking, however not screened a second time as it is discarded as duplicate. Additionally, further research on the characteristics and predictors of falsely excluded studies is needed. Thus, a methods study that analyzes a heterogeneous data set with a large quantity of screening decisions for possible predictors in a regression analysis or prediction model is warranted.

Conclusions

We cannot draw any firm conclusion about the most reliable and most valid method to recover studies falsely excluded during literature screening, as the available evidence is limited to a single-case study. Furthermore, due to the limited evidence from two case studies and one case series, we can draw no firm conclusions on characteristics that might predict a higher risk of false exclusion.

Abbreviations

CADTH: Canadian Agency for Drugs and Technologies in Health; CI: Confidence interval; CONSORT: Consolidated Standards of Reporting Trials; DART : Digital Access to Research Theses; IMRaD: Introduction, Materials and Methods, Results, and Discussion and Conclusion; KQ: Key question; MeSH: Medical Subject Headings; N: Number; NA: Not applicable; NNR: Numbers needed to read; NR: Not reported; OR: Odds ratio; PRESS: Peer review of the electronic search strategy; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; QUOROM: QUality Of Reporting Of Meta-analyses; RCT: Randomized controlled trial; SR: Systematic review; STARD: Standards for Reporting Diagnostic Accuracy.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13643-022-02109-w>.

Additional file 1. PRISMA checklist.

Additional file 2. Search strategy for Medline, Science Citation Index Expanded, Social Sciences Citation Index, Current Contents Connect, Embase, Epistemonikos.org, and Information Science & Technology Abstracts.

Additional file 3. Search strategy for similar articles searches in PubMed, Google Scholar and Scopus.

Additional file 4. Excluded studies at the full-text level.

Additional file 5. Risk of bias of included studies.

Acknowledgements

We would like to thank Angela Kaminski-Hartenthaler, who supported the abstract screening. In particular, we want to thank Jos Kleijnen for his input on the manuscript.

Authors' contributions

LA, AD, and GG developed the study concept. LA wrote the protocol. IK conducted the literature searches and similar article searches. LA, AD, GW, IS, and EP screened the references and extracted the data. LA conducted gray literature searches, reference list checking, and hand searches. GG provided methodological input throughout the study. LA, AD, and EP created the first draft of the manuscript, which GG, GW, IS, and IK critically revised. The authors read and approved the final version of the submitted manuscript.

Funding

This work was supported by a scholarship for the first author (LA) of the Gesellschaft für Forschungsförderung Niederösterreich m.b.H.

Availability of data and materials

All data has been summarized and provided in the manuscript or supplementary files.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department for Evidence-Based Medicine and Evaluation, Cochrane Austria, Danube University Krems, Dr. Karl Dorrek Strasse 30, 3500 Krems, Austria.

²Department of Family Medicine, Care and Public Health Research Institute (CAPHRI), Maastricht University, Peter Debyeplein 1, 6229 HA Maastricht, The Netherlands. ³RTI International, 3040 Cornwallis Road, PO Box 12194, Research Triangle Park, North Carolina 27709-2194, USA.

Received: 20 July 2021 Accepted: 30 October 2022

Published online: 09 November 2022

References

- Tufanaru C, Munn Z, Aromataris E, Campbell J, Hopp L. Chapter 3: Systematic reviews of effectiveness. *JBI Manual for Evidence Synthesis*. JBI, Adelaide; 2020.
- Centre for Reviews Dissemination. *CRD's guidance for undertaking reviews in healthcare*. York: York Publ. Services; 2009.
- Institute for Quality and Efficiency in Health. *General Methods Version 6.0 2020*. Köln: Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen; 2020.
- Shemilt I, Khan N, Park S, Thomas J. Use of cost-effectiveness analysis to compare the efficiency of study identification methods in systematic reviews. *Syst Rev*. 2016;5(1):140.
- Gartlehner G, Affengruber L, Titscher V, Noel-Storr A, Dooley G, Ballarini N, et al. Single-reviewer abstract screening missed 13 percent of relevant studies: a crowd-based, randomized controlled trial. *J Clin Epidemiol*. 2020;121:20–8.
- Waffenschmidt S, Knelangen M, Sieben W, Bühn S, Pieper D. Single screening versus conventional double screening for study selection in systematic reviews: a methodological systematic review. *BMC Med Res Methodol*. 2019;19(1):132.
- EUnetHTA JA3WP6B2-2 Authoring Team. Process of information retrieval for systematic reviews and health technology assessments on clinical effectiveness. *Methodological Guidelines*. Diemen (The Netherlands): EUnetHTA; 2019. Available from <https://www.eunetha.eu/>.
- Lefebvre C, Glanville J, Briscoe S, Littlewood A, Marshall C, Metzendorf MI, et al. Searching for and selecting studies. *Cochrane Handbook for Systematic Reviews of Interventions*. 2019:67–107.
- Tricco AC, Langlois E, Straus SE, Organization WH. *Rapid reviews to strengthen health policy and systems: a practical guide*. Geneva: World Health Organization; 2017.
- Horsley T, Dingwall O, Sampson M. Checking reference lists to find additional studies for systematic reviews. *Cochrane Database Syst Rev*. 2011;2011(8):MR000026.
- Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021)*. Cochrane, editor. 2021.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.
- Canadian Agency for Drugs and Technologies in Health (CADTH). *Grey matters: a practical tool for searching health-related grey literature*. 2019.
- Moola S, Z, Tufanaru C, Aromataris E, Sears K, Sfetcu R, Currie M, Lisy K, Qureshi R, Mattis P, Mu P. Chapter 7: systematic reviews of etiology and risk. *Adelaide: JBI*; 2020.
- Rathbone J, Albarqouni L, Bakhit M, Beller E, Byambasuren O, Hoffmann T, et al. Expediting citation screening using PICO-based title-only screening for identifying studies in scoping searches and rapid reviews. *Syst Rev*. 2017;6(1):1–7.
- Busse JW, Bruno P, Malik K, Connell G, Torrance D, Ngo T, et al. An efficient strategy allowed English-speaking reviewers to identify foreign-language articles eligible for a systematic review. *J Clin Epidemiol*. 2014;67(5):547–53.
- Feehan LM, Beck CA, Harris SR, MacIntyre DL, Li LC. Exercise prescription after fragility fracture in older adults: a scoping review. *Osteoporos Int*. 2011;22(5):1289–322.
- Beck C. The scoping study in physical therapy: application of traditional systematic review guidelines to an emerging methodology. Ontario: Canadian Health Libraries Association Conference in Kingston; 2010.
- Hamel C, Hersi M, Kelly SE, Tricco AC, Straus S, Wells G, et al. Guidance for using artificial intelligence for title and abstract screening while conducting knowledge syntheses. *BMC Med Res Methodol*. 2021;21(1):285.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

