

LETTER

Open Access



Enhanced recovery after surgery (ERAS) for vascular surgery: an evidence map and scoping review

Eric A. Apaydin^{1,2,3*} , Karen Woo⁴, Julia Rollison⁵, Sangita Baxi², Aneesha Motala^{1,2} and Susanne Hempel^{1,2}

Abstract

Background Enhanced recovery after surgery (ERAS) interventions aim to improve patient outcomes. Vascular surgery patients have unique requirements and it is unclear which ERAS interventions are supported by an evidence base.

Methods We conducted a scoping review to identify ERAS randomized controlled trials (RCTs) published in the biomedical or nursing literature. We assessed interventions for applicability to vascular surgery and differentiated interventions given at preadmission, preoperative, intraoperative, and postoperative surgery stages. We documented the research in an evidence map.

Results We identified 76 relevant RCTs. Interventions were mostly administered in preoperative (23 RCTs; 30%) or intraoperative surgery stages (35 RCTs; 46%). The majority of studies reported mortality outcomes (44 RCTs; 58%), but hospital (27 RCTs; 35%) and intensive care unit (9 RCTs; 12%) length of stay outcomes were less consistently described.

Conclusion The ERAS evidence base is growing but contains gaps. Research on preadmission interventions and more consistent reporting of key outcomes is needed.

Keywords Enhanced recovery, Surgery, Vascular, Evidence map

Introduction

Enhanced recovery after surgery (ERAS) consists of interventions designed to support patients recovering from surgery throughout the continuum of care [1]. Improved recovery after surgery benefits patients, practitioners, and healthcare delivery organizations [2]. However, little is known about the applicability ERAS interventions to vascular surgery [3]. Vascular surgery, which includes a wide range of treatments that encompass major open operations and less invasive endovascular procedures, may benefit from ERAS approaches.

To better understand this literature, we conducted a scoping review and created an evidence map of randomized controlled trials (RCTs) for ERAS interventions, mapped by surgical stage, that evaluate key

*Correspondence:

Eric A. Apaydin
eapaydin@usc.edu

¹ Southern California Evidence Review Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

² RAND Health Care, RAND Corporation, Santa Monica, CA, USA

³ Center for the Study of Healthcare Innovation, Implementation, and Policy, VA Greater Los Angeles Healthcare System, Los Angeles, CA, USA

⁴ Department of Surgery, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, USA

⁵ RAND Health Care, RAND Corporation, Arlington, VA, USA



This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2023. **Open**

Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Table 1 Evidence table

Citation	Stage	Mortality	LOS	ICU LOS
Ali 2007 [6]	Intraoperative	Measured	Measured	Measured
Baldwin 1994 [7]	Postoperative	Measured		
Barlow 1989 [8]	Preoperative			
Belch 1980 [9]	Preoperative	Measured		
Bender 1997 [10]	Preoperative	Measured		
Berlauk 1991 [11]	Preoperative	Measured		Measured
Bille-Brahe 1980 [12]	Postoperative			
Bode 1996 [13]	Intraoperative	Measured	Measured	Measured
Bohner 2002 [14]	Postoperative	Measured	Measured	Measured
Bolliger 2007 [15]	Preoperative	Measured		
Bonazzi 2002 [16]	Intraoperative	Measured	Measured	
Brady 2005 [17]	Intraoperative	Measured	Measured	
Christopherson 1996 [18]	Intraoperative	Measured		
Cook 1986 [19]	Intraoperative	Measured		
Dorman 1995 [20]	Intraoperative			
Durazzo 2004 [21]	Preoperative	Measured		
Earnshaw 1989 [22]	Postoperative			
Farkas 1993 [23]	Multi-stage	Measured		
Fleisher 2005 [24]	Preoperative	Measured		
Fleron 2003 [25]	Intraoperative	Measured		
Forster 2006 [26]	Postoperative			
Fourneau 2006 [27]	Intraoperative		Measured	
Frank 1992 [28]	Intraoperative			
Friedman 1996 [29]	Postoperative		Measured	Measured
Gonzalez-Fajardo 2009 [30]	Postoperative		Measured	
Gouaillier-Vulcain 2015 [31]	Intraoperative			
Hall 1998 [32]	Preoperative			
Hasselgren 1984 [33]	Preoperative			
Healy 2015 [34]	Intraoperative	Measured	Measured	Measured
Kaiser 1978 [35]	Intraoperative			
Kavakli 2019 [36]	Intraoperative		Measured	Measured
Kouvelos 2013 [37]	Preoperative			
Krog 2017 [38]	Intraoperative	Measured	Measured	
Kucukakin 2010 [39]	Intraoperative			
Kwon 2018 [40]	Postoperative		Measured	
Lee 2017 [41]	Postoperative	Measured	Measured	
Lindholm 2013 [42]	Intraoperative	Measured	Measured	
Linni 2012 [43]	Preoperative	Measured	Measured	
Lundorff 1999 [44]	Intraoperative			
Lunen 2018 [45]	Preoperative	Measured	Measured	Measured
Marroni 1999 [46]	Preoperative	Measured		
Martin 1982 [47]	Intraoperative			
Miller 1994 [48]	Postoperative	Measured	Measured	
Monsel 2016 [49]	Postoperative			
Mouren 1989 [50]	Intraoperative			
Muehling 2009 [51]	Multi-stage	Measured	Measured	Measured
Murphy 2014 [52]	Intraoperative	Measured	Measured	

Table 1 (continued)

Citation	Stage	Mortality	LOS	ICU LOS
Nesek-Adam 2012 [53]	Intraoperative			
Nevelsteen 1991 [54]	Preoperative			
Niemi 2006 [55]	Intraoperative			
Norgen 2004 [56]	Intraoperative	Measured		
Norris 2001 [57]	Multi-stage	Measured	Measured	
Oliver 2006 [58]	Intraoperative	Measured		
Ozaki 2015 [59]	Postoperative			
Partridge 2017 [60]	Preoperative	Measured	Measured	
Pitt 1980 [61]	Intraoperative	Measured		
Pleger 2018 [62]	Postoperative			Measured
Poldermans 1999 [63]	Intraoperative	Measured		
Reinhart 1989 [64]	Intraoperative	Measured		
Renghi 2013 [65]	Intraoperative			Measured
Risberg 1995 [66]	Preoperative	Measured		
Roizen 1980 [67]	Preoperative			
Salzmann 1983 [68]	Intraoperative			
Schouten 2009 [69]	Intraoperative	Measured		
Soliman 2016 [70]	Preoperative	Measured	Measured	
Sprung 2000 [71]	Intraoperative			
Stuhmeier 1996 [72]	Preoperative	Measured		
Subramaniam 2009 [73]	Intraoperative	Measured	Measured	
Swinnen 2010 [74]	Intraoperative			
Thomas 2016 [75]	Preoperative	Measured	Measured	
Turner 2008 [76]	Preoperative	Measured	Measured	Measured
Turtianien 2012 [77]	Postoperative	Measured	Measured	
Van der Linden 2010 [78]	Intraoperative	Measured		
Vierhout 2014 [79]	Postoperative			
Vukovic 2012 [80]	Postoperative			
Weed 2017 [81]	Preoperative			Measured
Worning 1986 [82]	Preoperative			
Yang 2006 [83]	Intraoperative	Measured	Measured	
Ziegler 1997 [84]	Preoperative	Measured		Measured

Abbreviations: ICU Intensive care unit, LOS Length of stay

patient-centered outcomes (mortality and length of stay). The evidence map [4] approach allows readers to view the existing evidence base in one figure and identify key research gaps.

Methods

As part of a larger project on ERAS, we searched PubMed and the Cumulative Index to Nursing and Allied Health Literature for published literature, and searched ClinicalTrials.gov for clinical trial records from inception to March 2023 (search strategies shown in Supplementary Material 1). Citations and full text publications were screened by experienced literature reviewers using




Fig. 1 Evidence map of reported outcomes in included ERAS studies by surgical stage

Abbreviations: *ICU* Intensive care unit, *LOS* Length of stay

predetermined eligibility criteria (full criteria shown in Supplementary Material 2). Notably, we restricted to RCTs, a study design that allows strong evidence statements. Abstracted items are also listed in Supplementary Material 2. Results were plotted as an evidence map using data visualization software (R Studio; R Studio Public Benefit Corporation; Boston, MA). All data analyzed in this manuscript is available in Table 1. This study did not involve human subjects and was therefore exempt from RAND Human Subjects Protection Committee review. Our full report on ERAS interventions for vascular surgery, with a search from inception to July 2019, is available on the Patient-Centered Outcomes Research Institute (PCORI) website [5].

Results

The search identified 4,483 citations, and we included 79 RCTs of ERAS interventions for vascular surgery (evidence table: Table 1) [6–84].

Figure 1 documents the evidence base and distribution of evidence across treatment stages and treatment outcomes. Of the evaluated interventions, 24 (30%) were preoperative, 36 (46%) were intraoperative, 16 (20%) were postoperative, and 3 (4%) were multi-stage. We did not identify any preadmission interventions. Forty-four RCTs (56%) reported mortality outcomes (Fig. 1), and of these interventions, 16 (36%) were preoperative, 20 (45%) were intraoperative, 5 (11%) were postoperative, and 3 (7%) were multi-stage. Length of stay outcomes were reported in 30 RCTs (38%), which evaluated ERAS interventions administered preoperatively (7 RCTs; 23%), intraoperatively (13 RCTs; 43%), postoperatively (8 RCTs; 27%), and across multiple surgical stages (2 RCTs; 7%). Length of stay in intensive care units (ICUs) was measured least frequently, in only 11 RCTs (14%). Of interventions in these studies, 4 (36%) were preoperative, 4 (36%) were intraoperative, 2 (18%) were postoperative, and 1 (9%) was multi-stage.

Discussion

We identified a substantial body of evidence of ERAS interventions for vascular surgery. Our analysis utilized an evidence map to categorize the available research on RCTs reporting mortality and length of stay outcomes for these interventions by surgical stage, a strategy not previously employed by other reviews. The map shows that existing research has primarily addressed pre- or intraoperative, rather than preadmission, surgical stages. In addition, while more than half of studies reported on mortality, information on hospital or ICU length of stay remains sparse.

A recent review [85] of 19 RCTs and observational studies of ERAS interventions for vascular surgery found that ERAS interventions reduced length of stay by 3.5 days across five studies. The review did not report pooled effects for mortality outcomes. A comprehensive systematic review and meta-analysis of the effects of ERAS interventions in the extant literature is needed to better estimate treatment effects.

Additional primary research on ERAS interventions is also needed. The existing evidence base lacks studies of preadmission ERAS interventions, and mortality and length of stay outcomes were not consistently reported. Future primary research should aim to study preadmission interventions and consistently measure and report mortality and length of stay outcomes.

ERAS research is rapidly growing and an evidence base for vascular surgery is also emerging. Our evidence map clearly outlines research gaps, including a lack of research on ERAS interventions at all surgery stages and the sparseness of information on key outcomes. ERAS interventions hold promise to improve patient recovery after vascular surgery and further applications should be explored.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13643-023-02324-z>.

Additional file 1: Supplementary Material 1. Search strategies. **Supplementary Material 2.** Eligibility criteria and data abstraction methods.

Acknowledgements

We thank Katharine McGinagle, Thomas Concannon, Paul Koegel, Jeanne Ringel, Rachel Andricosky, Michelle Althuis, and William Lawrence for helpful comments, Sachi Yagyu for conducting literature searches, and Christine Chen for research support.

Authors' contributions

Drs. Eric Apaydin, Karen Woo, Julia Rollison, and Susanne Hempel contributed to the study conception and design. Screening and abstraction were performed by Ms. Sangita Baxi, Dr. Julia Rollison, and Dr. Eric Apaydin. Ms. Sangita Baxi created the evidence maps, and Dr. Eric Apaydin analyzed the data. Ms. Aneesa Motala managed the screening and abstraction databases, and the overall project. The first draft of the manuscript was written by Dr. Eric Apaydin and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding

The manuscript is based on work funded by the Patient-Centered Outcomes Research Institute (contract holder: SH; PCORI [www.pcori.org]; Analytic and Operational Support for the Engagement of the Public and Patients Task Order #18 Topic Briefs on Enhanced Recovery After Surgery and Infantile Epilepsy). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

All data analyzed in this manuscript is available in Table 1.

Declarations

Ethics approval and consent to participate

This study did not involve human subjects and was therefore exempt from RAND Human Subjects Protection Committee review. No human subjects were involved, and thus their consent to participate was not applicable.

Consent for publication

All authors have given consent for publication.

Competing interests

The authors have declared that no competing interests exist.

Received: 28 June 2023 Accepted: 15 August 2023

Published online: 14 September 2023

References

1. Melloul E, Hubner M, Scott M, et al. Guidelines for Perioperative Care for Liver Surgery: Enhanced Recovery After Surgery (ERAS) Society Recommendations. *World J Surg.* 2016;40(10):2425–40. <https://doi.org/10.1007/s00268-016-3700-1>.
2. Society for Vascular Surgery. What is a vascular surgeon?. <https://vascular.org/patient-resources/what-vascular-surgeon>. Accessed 29 Apr 2022.
3. Society for Vascular Surgery. Clinical Practice Guidelines. <https://vascular.org/research-quality/guidelines-and-reporting-standards/clinical-practice-guidelines>. Accessed 29 Apr 2022.
4. Miake-Lye IM, Hempel S, Shanman R, Shekelle PG. What is an evidence map? A systematic review of published evidence maps and their definitions, methods, and products. *Systematic reviews.* 2016;5:28. <https://doi.org/10.1186/s13643-016-0204-x>.
5. Rollison J, Woo K, Chen C, Yagyu S, Motala A, Hempel S. Enhanced Recovery After Surgery for Vascular Surgery. 2019. <https://www.pcori.org/sites/default/files/PCORI-Topic-Brief-Enhanced-Recovery-After-Surgery-for-Vascular-Surgery.pdf>.
6. Ali ZA, Callaghan CJ, Lim E, et al. Remote ischemic preconditioning reduces myocardial and renal injury after elective abdominal aortic aneurysm repair: a randomized controlled trial. *Circulation.* 2007;116(11 Suppl):198–105. Intraoperative.
7. Baldwin L, Henderson A, Hickman P. Effect of postoperative low-dose dopamine on renal function after elective major vascular surgery. *Ann Int Med.* 1994;120(9):744–7. Postoperative.
8. Barlow IW, Ausubsky JR, Wilkinson D, Kester RC. Controlled trial of cephadrine versus cefuroxime in vascular surgery. *Int J Clin Pharmacol Res.* 1989;9(3):223–7. Preoperative.
9. Belch JJ, Lowe GD, Pollock JG, Forbes CD, Prentice CR. Low dose heparin in the prevention of deep-vein thrombosis after aortic bifurcation graft surgery. *Thromb Haemost.* 1980;42(5):1429–33. Preoperative.
10. Bender JS, Smith-Meek MA, Jones CE. Routine pulmonary artery catheterization does not reduce morbidity and mortality of elective vascular surgery: results of a prospective, randomized trial. *Ann Surg.* 1997;226(3):229–36. discussion 236–7. Preoperative.
11. Berlauck JF, Abrams JH, Gilmour IJ, O'Connor SR, Knighton DR, Cerra FB. Preoperative optimization of cardiovascular hemodynamics improves outcome in peripheral vascular surgery. A prospective, randomized clinical trial. *Ann Surg.* 1991;214(3):289–97. discussion 298–9. Preoperative.
12. Bille-Brahe NE, Engell HC, Sorensen MB. Acute postoperative digitalization of patients with arteriosclerotic heart disease after major surgery. A randomized haemodynamic study and proposal for therapy. *Acta anaesthesiologica Scandinavica.* 1980;24(6):501–6. Postoperative.
13. Bode RH Jr, Lewis KP, Zarich SW, et al. Cardiac outcome after peripheral vascular surgery. Comparison of general and regional anesthesia. *Anesthesiology.* 1996;84(1):3–13. Intraoperative.
14. Bohner H, Kindgen-Milles D, Grust A, et al. Prophylactic nasal continuous positive airway pressure after major vascular surgery: results of a prospective randomized trial. *Langenbeck's Arch Surg.* 2002;387(1):21–6. Postoperative.
15. Bolliger D, Seeberger MD, Lurati Buse GA, Christen P, Gurke L, Filipovic M. Randomized clinical trial of moxonidine in patients undergoing major vascular surgery. *British J Surg.* 2007;94(12):1477–84. Preoperative.
16. Bonazzi M, Gentile F, Biasi GM, et al. Impact of perioperative haemodynamic monitoring on cardiac morbidity after major vascular surgery in low risk patients. A randomised pilot trial. *Eur J Vasc Endovasc Surg.* 2002;23(5):445–51. Intraoperative.
17. Brady AR, Gibbs JS, Greenhalgh RM, Powell JT, Sydes MR. Perioperative beta-blockade (POBBLE) for patients undergoing infrarenal vascular surgery: results of a randomized double-blind controlled trial. *J Vasc Surg.* 2005;41(4):602–9. Intraoperative.
18. Christopherson R, Glavan NJ, Norris EJ, et al. Control of blood pressure and heart rate in patients randomized to epidural or general anesthesia for lower extremity vascular surgery. Perioperative Ischemia Randomized Anesthesia Trial (PIRAT) Study Group. *J Clin Anesth.* 1996;8(7):578–84. Intraoperative.
19. Cook PT, Davies MJ, Cronin KD, Moran P. A prospective randomised trial comparing spinal anaesthesia using hyperbaric cinchocaine with general anaesthesia for lower limb vascular surgery. *Anaesth Intensive Care.* 1986;14(4):373–80. Intraoperative.
20. Dorman BH, Elliott BM, Spinale FG, et al. Protamine use during peripheral vascular surgery: a prospective randomized trial. *J Vasc Surg.* 1995;22(3):248–55. discussion 256. Intraoperative.
21. Durazzo AE, Machado FS, Ikeoka DT, et al. Reduction in cardiovascular events after vascular surgery with atorvastatin: a randomized trial. *J Vasc Surg.* 2004;39(5):967–75. discussion 975–6. Preoperative.
22. Earnshaw JJ, Berridge DC, Slack RC, Makin GS, Hopkinson BR. Do pre-operative chlorhexidine baths reduce the risk of infection after vascular reconstruction? *Eur J Vasc Surg.* 1989;3(4):323–6. Postoperative.
23. Farkas JC, Chapuis C, Combe S, et al. A randomised controlled trial of a low-molecular-weight heparin (Enoxaparin) to prevent deep-vein thrombosis in patients undergoing vascular surgery. *Eur J Vasc Surg.* 1993;7(5):554–60. Bundle/unclear.
24. Fleisher LA, Newman MF, St Aubin LB, et al. Efficacy of zonisporide, an Na/H exchange ion inhibitor, for reducing perioperative cardiovascular events in vascular surgery patients. *J Cardiothorac Vasc Anesth.* 2005;19(5):570–6. Preoperative.
25. Fleron MH, Weiskopf RB, Bertrand M, et al. A comparison of intrathecal opioid and intravenous analgesia for the incidence of cardiovascular, respiratory, and renal complications after abdominal aortic surgery. *Anesth Analg.* 2003;97(1):2–12. table of contents. Intraoperative.
26. Forster JG, Rosenberg PH, Niemi TT. Continuous spinal microcatheter (28 gauge) technique for arterial bypass surgery of the lower extremities and comparison of ropivacaine with or without morphine for postoperative analgesia. *British J Anaesth.* 2006;97(3):393–400. Postoperative.
27. Fourneau I, Sabbe T, Daenens K, Nevelsteen A. Hand-assisted laparoscopy versus conventional median laparotomy for aortobifemoral bypass for severe aorto-iliac occlusive disease: a prospective randomised study. *Eur J Vasc Endovasc Surg.* 2006;32(6):645–50. Intraoperative.
28. Frank SM, Beattie C, Christopherson R, et al. Epidural versus general anesthesia, ambient operating room temperature, and patient age as predictors of inadvertent hypothermia. *Anesthesiology.* 1992;77(2):252–7. Intraoperative.
29. Friedman SG, Sowerby SA, Del Pin CA, Scher LA, Tortolani AJ. A prospective randomized study of abdominal aortic surgery without postoperative nasogastric decompression. *Cardiovasc Surg (London, England).* 1996;4(4):492–4.
30. Gonzalez-Fajardo JA, Mengibar L, Brizuela JA, Castrodeza J, Vaquero-Puerta C. Effect of postoperative restrictive fluid therapy in the recovery of patients with abdominal vascular surgery. *Eur J Vasc Endovasc Surg.* 2009;37(5):538–43. Postoperative.
31. Gouaillier-Vulcain F, Marchand E, Martinez R, Picquet J, Enon B. Utility of electrofusion for the femoral approach in vascular surgery: a randomized prospective study. *Ann Vasc Surg.* 2015;29(4):801–9. Intraoperative.
32. Hall JC, Christiansen KJ, Goodman M, et al. Duration of antimicrobial prophylaxis in vascular surgery. *Am J Surg.* 1998;175(2):87–90. Preoperative.
33. Hasselgren PO, Ivarsson L, Risberg B, Seeman T. Effects of prophylactic antibiotics in vascular surgery. A prospective, randomized, double-blind study. *Ann Surg.* 1984;200(1):86–92. Preoperative.
34. Healy DA, Boyle E, McCartan D, et al. A multiCenter pilot randomized controlled trial of remote ischemic preconditioning in major vascular surgery. *Vasc Endovasc Surg.* 2015;49(8):220–7. Intraoperative.
35. Kaiser AB, Clayton KR, Mulherin JL Jr, et al. Antibiotic prophylaxis in vascular surgery. *Ann Surg.* 1978;188(3):283–9. Intraoperative.

36. Kavakli AS, Kavrut Ozturk N, Yavuzel Adas H, et al. The effects of music on anxiety and pain in patients during carotid endarterectomy under regional anesthesia: a randomized controlled trial. *Complement Ther Med*. 2019;44:94–101.
37. Kouvelos GN, Arnaoutoglou EM, Matsagkas MI, et al. Effects of rosuvastatin with or without ezetimibe on clinical outcomes in patients undergoing elective vascular surgery: results of a pilot study. *J Cardiovasc Pharmacol Ther*. 2013;18(1):5–12. Preoperative.
38. Krog AH, Thorsby PM, Sahba M, et al. Perioperative humoral stress response to laparoscopic versus open aortobifemoral bypass surgery. *Scand J Clin Lab Investigation*. 2017;77(2):83–92. Intraoperative.
39. Kucukakin B, Wilhelmsen M, Lykkesfeldt J, Reiter RJ, Rosenberg J, Gogenuur I. No effect of melatonin to modify surgical-stress response after major vascular surgery: a randomised placebo-controlled trial. *Eur J Vasc Endovasc Surg*. 2010;40(4):461–7. Intraoperative.
40. Kwon J, Staley C, McCullough M, et al. A randomized clinical trial evaluating negative pressure therapy to decrease vascular groin incision complications. *J Vasc Surg*. 2018;68(6):1744–52. Postoperative.
41. Lee K, Murphy PB, Ingves MV, et al. Randomized clinical trial of negative pressure wound therapy for high-risk groin wounds in lower extremity revascularization. *J Vasc Surg*. 2017;66(6):1814–9. Postoperative.
42. Lindholm EE, Aune E, Noren CB, et al. The anesthesia in abdominal aortic surgery (ABSENT) study: a prospective, randomized, controlled trial comparing rocuronium T release with fentanyl-sevoflurane and propofol-remifentanyl anesthesia in major vascular surgery. *Anesthesiology*. 2013;119(4):802–12. Intraoperative.
43. Linni K, Mader N, Aspalter M, et al. Ultrasonic vein mapping prior to infrainguinal autogenous bypass grafting reduces postoperative infections and readmissions. *J Vasc Surg*. 2012;56(1):126–32. discussion 132–3. Preoperative.
44. Lunderhoff L, Dich-Nielsen JO, Laugesen H, Jensen MM. Single-dose spinal anaesthesia versus incremental dosing for lower limb vascular surgery. *Acta Anaesth Scand*. 1999;43(4):405–10. Intraoperative.
45. Lunen TB, Johansson PI, Jensen LP, et al. Administration of platelets to ruptured abdominal aortic aneurysm patients before open surgery: a prospective, single-blinded, randomised study. *Transfusion Med (Oxford, England)*. 2018;28(5):386–91. Preoperative.
46. Marroni M, Cao P, Fiorio M, et al. Prospective, randomized, double-blind trial comparing teicoplanin and cefazolin as antibiotic prophylaxis in prosthetic vascular surgery. *Eur J Clin Microbiol Infect Dis*. 1999;18(3):175–8. Preoperative.
47. Martin DE, Rosenberg H, Aukburg SJ, et al. Low-dose fentanyl blunts circulatory responses to tracheal intubation. *Anesth Analg*. 1982;61(8):680–4. Intraoperative.
48. Miller DR, Martineau RJ, Hull KA, Vallee F, LeBel M. Optimizing sedation following major vascular surgery: a double-blind study of midazolam administered by continuous infusion. *Can J Anaesth*. 1994;41(9):782–93. Postoperative.
49. Monsel A, Lu Q, Le Corre M, et al. Tapered-cuff endotracheal tube does not prevent early postoperative pneumonia compared with spherical-cuff endotracheal tube after major vascular surgery: a randomized controlled trial. *Anesthesiology*. 2016;124(5):1041–52. Postoperative.
50. Mouren S, Baron JF, Hag B, Arthaud M, Vials P. Normovolemic hemodilution and lumbar epidural anesthesia. *Anesth Analg*. 1989;69(2):174–9. Intraoperative.
51. Muehling B, Schelzig H, Steffen P, Meierhenrich R, Sunder-Plassmann L, Orend KH. A prospective randomized trial comparing traditional and fast-track patient care in elective open infrarenal aneurysm repair. *World J Surg*. 2009;33(3):577–85. Bundle/unclear.
52. Murphy N, Vijayan A, Frohlich S, et al. Remote ischemic preconditioning does not affect the incidence of acute kidney injury after elective abdominal aortic aneurysm repair. *J Cardiothoracic Vasc Anesth*. 2014;28(5):1285–92. Intraoperative.
53. Nesek-Adam V, Rasic Z, Schwarz D, et al. The effect of spinal versus general anesthesia on postoperative pain and analgesic requirements in patients undergoing peripheral vascular surgery. *Coll Antropol*. 2012;36(4):1301–5. Intraoperative.
54. Nevelsteen A, Mortelmans L, Van de Cruys A, Merckx E, Verhaeghe R. Effect of ticlopidine on blood loss, platelet turnover and platelet deposition on prosthetic surfaces in patients undergoing aorto-femoral bypass grafting. *Thromb Res*. 1991;64(3):363–9. Preoperative.
55. Niemi TT, Munsterhjelm E, Poyhia R, Hynninen MS, Salmenpera MT. The effect of N-acetylcysteine on blood coagulation and platelet function in patients undergoing open repair of abdominal aortic aneurysm. *Blood Coagul Fibrinolysis*. 2006;17(1):29–34. Intraoperative.
56. Norgren L. Can low molecular weight heparin replace unfractionated heparin during peripheral arterial reconstruction? An open label prospective randomized controlled trial. *J Vasc Surg*. 2004;39(5):977–84. Intraoperative.
57. Norris EJ, Beattie C, Perler BA, et al. Double-masked randomized trial comparing alternate combinations of intraoperative anesthesia and postoperative analgesia in abdominal aortic surgery. *Anesthesiology*. 2001;95(5):1054–67. Bundle/Mixed (multiple arms).
58. Oliver WC Jr, Nuttall GA, Cherry KJ, Decker PA, Bower T, Ereth MH. A comparison of fenoldopam with dopamine and sodium nitroprusside in patients undergoing cross-clamping of the abdominal aorta. *Anesth Analg*. 2006;103(4):833–40. Intraoperative.
59. Ozaki CK, Hamdan AD, Barshes NR, et al. Prospective, randomized, multi-institutional clinical trial of a silver alginate dressing to reduce lower extremity vascular surgery wound complications. *J Vasc Surg*. 2015;61(2):419–27.e1. Postoperative.
60. Partridge JS, Harari D, Martin FC, et al. Randomized clinical trial of comprehensive geriatric assessment and optimization in vascular surgery. *British J Surg*. 2017;104(6):679–87. Preoperative.
61. Pitt HA, Postier RG, MacGowan AW, et al. Prophylactic antibiotics in vascular surgery. Topical, systemic, or both? *Ann Surg*. 1980;192(3):356–64. Intraoperative.
62. Plegier SP, Nink N, Elzien M, Kunold A, Koshty A, Boning A. Reduction of groin wound complications in vascular surgery patients using closed incision negative pressure therapy (ciNPT): a prospective, randomised, single-institution study. *Int J Wound J*. 2018;15(1):75–83. Postoperative.
63. Poldermans D, Boersma E, Bax JJ, et al. The effect of bisoprolol on perioperative mortality and myocardial infarction in high-risk patients undergoing vascular surgery. Dutch echocardiographic cardiac risk evaluation applying Stress echocardiography study Group. *N Engl J Med*. 1999;341(24):1789–94. Intraoperative.
64. Reinhart K, Foehring U, Kersting T, et al. Effects of thoracic epidural anesthesia on systemic hemodynamic function and systemic oxygen supply-demand relationship. *Anesth Analg*. 1989;69(3):360–9. Intraoperative.
65. Renghi A, Gramaglia L, Casella F, Moniaci D, Gaboli K, Brustia P. Local versus epidural anesthesia in fast-track abdominal aortic surgery. *J Cardiothorac Vasc Anesth*. 2013;27(3):451–8. Intraoperative.
66. Risberg B, Drott C, Dalman P, et al. Oral ciprofloxacin versus intravenous cefuroxime as prophylaxis against postoperative infection in vascular surgery: a randomised double-blind, prospective multicentre study. *Eur J Vasc Endovasc Surg*. 1995;10(3):346–51. Preoperative.
67. Roizen MF, Sohn YJ, L'Hommedieu CS, Wylie EJ, Ota MK. Operating room temperature prior to surgical draping: effect on patient temperature in recovery room. *Anesth Analg*. 1980;59(11):852–5. Preoperative.
68. Salzmann G. Perioperative infection prophylaxis in vascular surgery—a randomized prospective study. *Thoracic Cardiovasc Surgeon*. 1983;31(4):239–42. Intraoperative.
69. Schouten O, Boersma E, Hoeks SE, et al. Fluvastatin and perioperative events in patients undergoing vascular surgery. *New England J Med*. 2009;361(10):980–9. Intraoperative.
70. Soliman R, Zohry G. The myocardial protective effect of dexmedetomidine in high-risk patients undergoing aortic vascular surgery. *Ann Cardiac Anaesth*. 2016;19(4):606–13. Preoperative.
71. Sprung J, Bourke DL, Schoenwald PK, et al. Small-dose dopamine increases epidural lidocaine requirements during peripheral vascular surgery in elderly patients. *Anesth Analg*. 2000;90(2):388–92. Intraoperative.
72. Stuhmeier KD, Mainzer B, Cierpka J, Sandmann W, Tarnow J. Small, oral dose of clonidine reduces the incidence of intraoperative myocardial ischemia in patients having vascular surgery. *Anesthesiology*. 1996;85(4):706–12. Preoperative.
73. Subramaniam B, Panzica PJ, Novack V, et al. Continuous perioperative insulin infusion decreases major cardiovascular events in patients undergoing vascular surgery: a prospective, randomized trial. *Anesthesiology*. 2009;110(5):970–7. Intraoperative.
74. Swinnen J, Chao A, Tiwari A, Crozier J, Vicaretti M, Fletcher J. Vertical or transverse incisions for access to the femoral artery: a randomized control study. *Ann Vasc Surg*. 2010;24(3):336–41. Intraoperative.

75. Thomas KN, Cotter JD, Williams MJ, van Rij AM. Repeated episodes of remote ischemic preconditioning for the prevention of myocardial injury in vascular surgery. *Vasc Endovascular Surg.* 2016;50(3):140–6. Preoperative.
76. Turner S, Derham C, Orsi NM, Bosomworth M, Bellamy MC, Howell SJ. Randomized clinical trial of the effects of methylprednisolone on renal function after major vascular surgery. *British J Surg.* 2008;95(1):50–6. Preoperative.
77. Turtiainen J, Saimanen EI, Makinen KT, et al. Effect of triclosan-coated sutures on the incidence of surgical wound infection after lower limb revascularization surgery: a randomized controlled trial. *World J Surg.* 2012;36(10):2528–34. Postoperative.
78. Van der Linden PJ, Dierick A, Wilmin S, Bellens B, De Hert SG. A randomized controlled trial comparing an intraoperative goal-directed strategy with routine clinical practice in patients undergoing peripheral arterial surgery. *Eur J Anaesthesiol.* 2010;27(9):788–93. Intraoperative.
79. Vierhout BP, Ott A, Reijnen MM, et al. Cyanoacrylate skin microsealant for preventing surgical site infection after vascular surgery: a discontinued randomized clinical trial. *Surg Infections.* 2014;15(4):425–30. Postoperative.
80. Vukovic J, Ramakrishnan P, Milan Z. Does epidural clonidine improve postoperative analgesia in major vascular surgery? *Med Glas Ljek komore Zenicko-dobojsk kantona.* 2012;9(1):49–55. Postoperative.
81. Weed HG. 2017 - Preoperative geriatric assessment reduced hospital stay and complications in patients ≥ 65 y having vascular surgery. *ACP J Club.* 2017;167(2):1.
82. Worning AM, Frimodt-Moller N, Ostri P, Nilsson T, Hojholdt K, Frimodt-Moller C. Antibiotic prophylaxis in vascular reconstructive surgery: a double-blind placebo-controlled study. *J Antimicrob Chemother.* 1986;17(1):105–13. Preoperative.
83. Yang H, Raymer K, Butler R, Parlow J, Roberts R. The effects of perioperative beta-blockade: results of the Metoprolol after Vascular Surgery (MaVS) study, a randomized controlled trial. *Am heart J.* 2006;152(5):983–90. Intraoperative.
84. Ziegler DW, Wright JG, Choban PS, Flancbaum L. A prospective randomized trial of preoperative "optimization" of cardiac function in patients undergoing elective peripheral vascular surgery. *Surgery.* 1997;122(3):584–92. Preoperative.
85. McGinagle KL, Eldrup-Jorgensen J, McCall R, et al. A systematic review of enhanced recovery after surgery for vascular operations. *J Vasc Surg.* 2019;70(2):629–40. <https://doi.org/10.1016/j.jvs.2019.01.050>. e1.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

